


# Quantum Optics and Information

Alexey V. Gorshkov

Joint Quantum Institute (JQI)

Joint Center for Quantum Information and Computer Science (QIACS)

NIST and University of Maryland



JOINT CENTER FOR  
QUANTUM INFORMATION  
AND COMPUTER SCIENCE

NIST



2021 Boulder Summer School

July 19, 20, 21

2021

# Outline for the three lectures

- interacting photons in Rydberg media
- (• dynamics of quantum systems with long-range interactions)

# Interacting photons in Rydberg media

References:

Homework: check these out,  
especially the last two

Quantum optics:

- lecture notes for Misha Lukin's class Modern Atomic and Optical Physics II, compiled by Lily Childress: <https://lukin.physics.harvard.edu/teaching>
- Meystre and Sargent, "Elements of Quantum Optics"
- Loudon, "Quantum Theory of Light"

Waveguide QED:

- Roy, Wilson, Firstenberg, "Colloquium: Strongly interacting photons in one-dimensional continuum," RMP 89, 021001 (2017)

Interacting photons in Rydberg media:

- Murray, Pohl, "Quantum and Nonlinear Optics in Strongly Interacting Atomic Ensembles", Adv. At., Mol., Opt. Phys. 65, 321 (2016)

Thanks to colleagues whose slides I am borrowing:  
Misha Lukin, Bill Phillips, Thomas Pohl,...

# Outline

- motivation and basic idea
- E&M field quantization
- propagation of light through atomic ensembles; electromagnetically induced transparency (EIT)
- Rydberg atoms
- basic idea revisited
- photon interacting with stationary excitation
  - on resonance: single-photon switch, subtractor
  - off resonance: two-photon quantum gate
- dynamics of multiple photons
  - on resonance: source of single photons
  - off resonance: two-photon gate, bound states, many-body physics
- more applications



# Outline

- motivation and basic idea
- E&M field quantization
- propagation of light through atomic ensembles; electromagnetically induced transparency (EIT)
- Rydberg atoms
- basic idea revisited
- photon interacting with stationary excitation
  - on resonance: single-photon switch, subtractor
  - off resonance: two-photon quantum gate
- dynamics of multiple photons
  - on resonance: source of single photons
  - off resonance: two-photon gate, bound states, many-body physics
- more applications

# Photons interact only in Sci-Fi




But light CAN act on light  
under the right circumstances





One of the first scientific achievements  
following the first laser...



...was the first demonstration of non-linear optics

VOLUME 7, NUMBER 4

PHYSICAL REVIEW LETTERS

AUGUST 15, 1961

---


GENERATION OF OPTICAL HARMONICS\*

P. A. Franken, A. E. Hill, C. W. Peters, and G. Weinreich


The Harrison M. Randall Laboratory of Physics, The University of Michigan, Ann Arbor, Michigan

(Received July 21, 1961)

# The first non-linear optics needed the first laser



Editor re-touched **signal** out!




# Electrons vs. Photons



$$F = \frac{1}{4\pi\epsilon_0} \frac{e^2}{r^2} \quad (\text{large})$$

transistor




small electrical signals  
control huge currents



photons hardly interact

nonlinear  
crystal



even huge optical intensities  
only control tiny optical signals




# Electronics vs. Photonics

electrons  
process information



photons  
transmit information






# Electronics vs. Photonics

electrons  
process information

photons  
transmit information



converting photons to electrical signals is “expensive”



# Electronics vs. Photonics

electrons  
process information



photons  
transmit information




Can we make photons process  
without conversion?

- photon-photon interactions too weak for processing information


# Photon-photon interactions

Typical approach to achieving interactions between optical photons:

- nonlinearity induced by individual atoms (or artificial atoms)



Kimble @ Caltech



Vuckovic @ Stanford

Hard!


# Photon-photon interactions

Typical approach to achieving interactions between optical photons:


- nonlinearity induced by individual atoms (or artificial atoms)

This talk:      Map strong **atom-atom interactions** onto  
strong photon-photon interactions

# Medium where photons interact strongly



Map strong **atom-atom interactions** onto  
**strong photon-photon interactions**



EIT = electromagnetically induced transparency

Experiments: Adams, Kuzmich, Lukin & Vuletic, Pfau & Löw, Grangier, Weidemüller, Hofferberth, Dürr & Rempe, Simon, Firstenberg, Ourjoumtsev, H. de Riedmatten, etc...  
Theory: Kurizki, Fleischhauer, Petrosyan, Mølmer, Pohl, Lesanovsky, Kennedy, Brion, Büchler, Sørensen, most experimental groups above, etc...

# Outline


- motivation and basic idea
- **E&M field quantization**
- propagation of light through atomic ensembles;  
electromagnetically induced transparency (EIT)
- Rydberg atoms
- basic idea revisited
- photon interacting with stationary excitation
  - on resonance: single-photon switch, subtractor
  - off resonance: two-photon quantum gate
- dynamics of multiple photons
  - on resonance: source of single photons
  - off resonance: two-photon gate, bound states,  
many-body physics
- more applications

# E&M field quantization

- Lukin/Childress lecture notes
- Meystre and Sargent, "Elements of Quantum Optics"
- consider free field (no sources)
- Maxwell's equations  $\Rightarrow$  wave equation

$$\nabla^2 \mathbf{E} - \frac{1}{c^2} \frac{\partial^2 \mathbf{E}}{\partial t^2} = 0$$

- knowing  $\mathbf{E}$ , find  $\mathbf{B}$  via  $\nabla \times \mathbf{B} = \frac{1}{c^2} \frac{\partial \mathbf{E}}{\partial t}$  (SI units used)



- large cavity of length  $L$  & volume  $V$ , with  $\mathbf{E} = 0$  on mirrors
- eigenmodes = standing waves

$$\sin(k_j z) \quad k_j L = \pi j$$

$$j = 1, 2, 3, \dots$$

( • could do periodic boundary conditions => running waves )

- consider  $\hat{x}$ -polarized field

$$E_x(z, t) = \sum_j \underbrace{\sqrt{\frac{2\nu_j^2}{\epsilon_0 V}}}_{A_j} \overset{\substack{\uparrow \\ \text{amplitude}}}{q_j(t)} \sin k_j z \quad \nu_j = ck_j$$

$$\Rightarrow B_y(z, t) = \frac{1}{c^2} \sum_j \frac{\dot{q}_j(t)}{k_j} A_j \cos k_j z$$



- classical energy:

$$H = \frac{1}{2} \int dV \left( \epsilon_0 E^2 + \frac{1}{\mu_0} B^2 \right)$$

$$= \sum_j \frac{\nu_j^2 q_j^2}{2} + \frac{\dot{q}_j^2}{2} \rightarrow \frac{1}{2} \sum_j (\nu_j^2 \hat{q}_j^2 + \hat{p}_j^2) = \sum_j \hbar \nu_j \left( \hat{a}_j^\dagger \hat{a}_j + \frac{1}{2} \right)$$

- independent harmonic oscillators with frequency  $\nu_j$ , unit mass, position  $q_j$

- Quantization:  $q_j \rightarrow \hat{q}_j$     $\dot{q}_j \rightarrow \hat{p}_j$     $[\hat{q}_j, \hat{p}_{j'}] = i\hbar \delta_{j,j'}$

- creation/annihilation:  $[\hat{a}_j, \hat{a}_{j'}^\dagger] = \delta_{j,j'}$

$$\hat{a}_j = \frac{1}{\sqrt{2\hbar\nu_j}} (\nu_j \hat{q}_j + i\hat{p}_j) \quad \hat{q}_j = \sqrt{\frac{\hbar}{2\nu_j}} (\hat{a}_j + \hat{a}_j^\dagger)$$

$$\hat{a}_j^\dagger = \frac{1}{\sqrt{2\hbar\nu_j}} (\nu_j \hat{q}_j - i\hat{p}_j) \quad \hat{p}_j = -i\sqrt{\frac{\hbar\nu_j}{2}} (\hat{a}_j - \hat{a}_j^\dagger)$$

- $\hat{x}$  component of electric field operator

$$\hat{E}_x(z) = \sum_j A_j \underbrace{\sqrt{\frac{\hbar}{2\nu_j}}}_{\sqrt{\frac{\hbar\nu_j}{\epsilon_0 V}} = \text{electric field per photon}} (\hat{a}_j + \hat{a}_j^\dagger) \sin k_j z$$

(makes sense:  $\hbar\nu_j \sim \text{energy} \sim \epsilon_0 E^2 V$  )

- for running waves, including all polarizations & directions

$$\hat{\mathbf{E}}(\mathbf{r}) = \hat{\mathcal{E}}(\mathbf{r}) + \hat{\mathcal{E}}^\dagger(\mathbf{r})$$

$$\hat{\mathcal{E}}(\mathbf{r}) = \sum_{\mathbf{k}, \alpha} \epsilon_\alpha \sqrt{\frac{\hbar\nu_j}{2\epsilon_0 V}} \hat{a}_{\mathbf{k}, \alpha} e^{i\mathbf{k} \cdot \mathbf{r}}$$

$\uparrow$   
 transverse  
 polarization

# Atom-field interactions

- starting point: dipole Hamiltonian for a 2-level atom

$$\hat{V}_{af} = -\hat{\mathbf{E}} \cdot \hat{\mathbf{d}} \quad |2\rangle \text{ —}$$

$$= -(\hat{\mathcal{E}} + \hat{\mathcal{E}}^\dagger) \cdot (\langle 2|\hat{\mathbf{d}}|1\rangle|2\rangle\langle 1| + \langle 1|\hat{\mathbf{d}}|2\rangle|1\rangle\langle 2|)$$

- 4 types of terms:

$|1\rangle \text{ —}$

$$\hat{a}|2\rangle\langle 1| \quad \cancel{\hat{a}|1\rangle\langle 2|} \quad \cancel{\hat{a}^\dagger|2\rangle\langle 1|} \quad \hat{a}^\dagger|1\rangle\langle 2|$$

(Heisenberg evolution under  $\hat{H} = \hbar\nu_j \hat{a}_j^\dagger \hat{a}_j$  :  $\hat{a}_j(t) = \hat{a}_j(0)e^{-i\nu_j t}$  )

- RWA  $\simeq$  energy conservation

- with RWA:  $\hat{V}_{af} = - \sum_{\mathbf{k}, \alpha} \hbar g_{\mathbf{k}, \alpha} |2\rangle\langle 1| \hat{a}_{j, \alpha} + \hbar g_{\mathbf{k}, \alpha}^* |1\rangle\langle 2| \hat{a}_{\mathbf{k}, \alpha}^\dagger$


- single-photon Rabi frequency:  $g_{\mathbf{k}, \alpha} = \frac{\mu_\alpha}{\hbar} \sqrt{\frac{\hbar\nu_j}{2\epsilon_0 V}} e^{i\mathbf{k} \cdot \mathbf{r}}$

- (if standing wave mode,  $|g| \propto \sin(kz)$  )  $\mu_\alpha = \langle 2|d_\alpha|1\rangle$

## Remarks

- no sources ( $\nabla \cdot \mathbf{E} = 0$ ), wave equation
    - $\Rightarrow$  didn't need  $\mathbf{A}$ ; used  $\hat{V}_{af} = -\hat{\mathbf{E}} \cdot \hat{\mathbf{d}}$
    - $\Rightarrow$  didn't need to choose gauge
  - with sources ( $\nabla \cdot \mathbf{E} \neq 0$ ), no wave equation
    - $\Rightarrow$  need  $\mathbf{A}$
    - $\Rightarrow$  choose Coulomb gauge  $\nabla \cdot \mathbf{A} = 0$
    - $\Rightarrow$  quantized similarly to this lecture
- [see Cohen-Tannoudji et al., “Photons and Atoms”]

# Medium where photons interact strongly





EIT = electromagnetically induced transparency

# Outline

- motivation and basic idea
- E&M field quantization
- **propagation of light through atomic ensembles;  
electromagnetically induced transparency (EIT)**
- Rydberg atoms
- basic idea revisited
- photon interacting with stationary excitation
  - on resonance: single-photon switch, subtractor
  - off resonance: two-photon quantum gate
- dynamics of multiple photons
  - on resonance: source of single photons
  - off resonance: two-photon gate, bound states,  
many-body physics
- more applications

# Three-level medium



$A$  = cross section of beam and of ensemble

$$\omega_1 = \omega_{eg} + \Delta$$

$$\omega_2 = \omega_{re} - \Delta$$

$$\hat{\mathbf{E}}_1(z) = \epsilon_1 \left( \frac{\hbar \omega_1}{4\pi c \epsilon_0 A} \right)^{1/2} \int d\omega \hat{a}_\omega e^{i\omega z/c} + h.c. \quad \text{Loudon, "Quantum Theory of Light"}$$



$$[\hat{a}_\omega, \hat{a}_{\omega'}^\dagger] = \delta(\omega - \omega')$$

$$\mathbf{E}_2(t) = \epsilon_2 \mathcal{E}_2(t) \cos(\omega_2 t)$$

Fleischhauer, Lukin, PRA 65, 022314 (2002)

AVG, Adre, Lukin, Sorensen, PRA 76, 033805 (2007)

# Three-level medium



$A$  = cross section of beam and of ensemble

$$\omega_1 = \omega_{eg} + \Delta$$

$$\omega_2 = \omega_{re} - \Delta$$

$$\hat{\mathbf{E}}_1(z) = \epsilon_1 \left( \frac{\hbar \omega_1}{4\pi c \epsilon_0 A} \right)^{1/2} \int d\omega \hat{a}_\omega e^{i\omega z/c} + h.c. \quad \text{Loudon, "Quantum Theory of Light"}$$

$$[\hat{a}_\omega, \hat{a}_{\omega'}^\dagger] = \delta(\omega - \omega')$$


$$\mathbf{E}_2(t) = \epsilon_2 \mathcal{E}_2(t) \cos(\omega_2 t)$$

Fleischhauer, Lukin, PRA 65, 022314 (2002)

AVG, Adre, Lukin, Sorensen, PRA 76, 033805 (2007)



# Three-level medium



$$\hat{H} = \hat{H}_0 + \hat{V}$$

$$\hat{H}_0 = \int d\omega \hbar \omega \hat{a}_\omega^\dagger \hat{a}_\omega + \sum_{i=1}^N \left( \hbar \omega_{rg} \hat{\sigma}_{rr}^i + \hbar \omega_{eg} \hat{\sigma}_{ee}^i \right)$$

$$\hat{V} = - \sum_{i=1}^N \hat{\mathbf{d}}_i \cdot [\mathbf{E}_2(t) + \hat{\mathbf{E}}_1(z_i)]$$

$$= - \hbar \sum_{i=1}^N \left( \Omega(t) \hat{\sigma}_{re}^i e^{-i\omega_2 t} + g \sqrt{\frac{1}{2\pi c}} \int d\omega \hat{a}_\omega e^{i\omega z_i/c} \hat{\sigma}_{eg}^i + h.c. \right)$$

$$\hat{\sigma}_{\mu\nu}^i = |\mu\rangle_{ii} \langle \nu|$$



$$\Omega(t) = \langle r | (\hat{\mathbf{d}} \cdot \epsilon_2) | e \rangle \mathcal{E}_2(t) / (2\hbar) = \text{Rabi frequency}$$

$$g = \langle e | (\hat{\mathbf{d}} \cdot \epsilon_1) | g \rangle \sqrt{\frac{\omega_1}{2\hbar\epsilon_0 A}}$$

- $\pi$  pulse takes time  $\pi/(2\Omega)$
- I will often set  $\hbar = 1$

# Three-level medium

define slowly varying operators



- thin enough that fields continuous
- assume almost all atoms in ground state at all times
- $n = \text{atom density}$

homework exercise

$$\hat{P}^\dagger(z, t) = \sqrt{n} \frac{1}{N_z} \sum_{i=1}^{N_z} \hat{\sigma}_{eg}^i(t) e^{-i\omega_1(t-z_i/c)}$$

creates  $|e\rangle$  excitation at  $z$

$$[\hat{P}(z, t), \hat{P}^\dagger(z', t)] = \delta(z - z')$$

$$\hat{S}^\dagger(z, t) = \sqrt{n} \frac{1}{N_z} \sum_{i=1}^{N_z} \hat{\sigma}_{rg}^i(t) e^{-i\omega_1(t-z_i/c) - i\omega_2 t}$$

creates  $|r\rangle$  excitation at  $z$


$$[\hat{S}(z, t), \hat{S}^\dagger(z', t)] = \delta(z - z')$$

$$\hat{\mathcal{E}}^\dagger(z, t) = \sqrt{\frac{1}{2\pi c}} e^{-i\omega_1(t-z/c)} \int d\omega \hat{a}_\omega^\dagger(t) e^{-i\omega z/c}$$

creates photon at  $z$

$$[\hat{\mathcal{E}}(z, t), \hat{\mathcal{E}}^\dagger(z', t)] = \delta(z - z')$$

# Three-level medium



$$\frac{\hat{H}}{\hbar} = -ic \int dz \hat{\mathcal{E}}^\dagger(z) \frac{\partial}{\partial z} \hat{\mathcal{E}}(z)$$

$$+ \int dz \left[ -\Delta \hat{P}^\dagger(z) \hat{P}(z) - \left( \Omega(t) \hat{S}^\dagger(z) \hat{P}(z) + g\sqrt{n} \hat{P}^\dagger(z) \hat{\mathcal{E}}(z) + h.c. \right) \right]$$

Heisenberg evolution:

$$(\partial_t + c\partial_z) \hat{\mathcal{E}} = ig\sqrt{n} \hat{P}$$

$$\partial_t \hat{P} = i\Delta \hat{P} + ig\sqrt{n} \hat{\mathcal{E}} + i\Omega \hat{S} - \gamma \hat{P} + \sqrt{2\gamma} \hat{F}_P$$

$$\partial_t \hat{S} = i\Omega^* \hat{P} - \gamma_s \hat{S} + \sqrt{2\gamma_s} \hat{F}_S$$

collective  
enhancement

Langevin noise


only nonzero noise correlations are:

$$\langle \hat{F}_P(z, t) \hat{F}_P^\dagger(z', t') \rangle = \delta(z - z') \delta(t - t')$$

$$\langle \hat{F} \rangle = \langle \hat{F} \hat{F} \rangle = \langle \hat{F}^\dagger \hat{F} \rangle = 0$$

$$\langle \hat{F}_S(z, t) \hat{F}_S^\dagger(z', t') \rangle = \delta(z - z') \delta(t - t')$$

# Three-level medium



$$(\partial_t + c\partial_z)\hat{\mathcal{E}} = ig\sqrt{n}\hat{P}$$

$$\partial_t\hat{P} = -(\gamma - i\Delta)\hat{P} + ig\sqrt{n}\hat{\mathcal{E}} + i\Omega\hat{S} + \sqrt{2\gamma}\hat{F}_P$$

$$\partial_t\hat{S} = -\gamma_s\hat{S} + i\Omega^*\hat{P} + \sqrt{2\gamma_s}\hat{F}_S$$

- assume all atoms initially in ground state, i.e. no P or S excitations
- assume 1 incoming photon

$$|\psi(t)\rangle = \int dz E(z, t) \hat{\mathcal{E}}^\dagger(z) |0\rangle + \int dz P(z, t) \hat{P}^\dagger(z) |0\rangle + \int dz S(z, t) \hat{S}^\dagger(z) |0\rangle$$

$$(\partial_t + c\partial_z)E = ig\sqrt{n}P$$

$$E(z, t = 0) = P(z, t = 0) = S(z, t = 0) = 0$$


$$\partial_t P = -(\gamma - i\Delta)P + ig\sqrt{n}E + i\Omega S$$

$$E(z = 0, t) = E_{in}(t)$$

$$\partial_t S = -\cancel{\gamma_s} S + i\Omega^* P$$

- same as equations for coherent input

# No atoms



$$(\partial_t + c\partial_z)E = ig\sqrt{n}P$$

$$\partial_t P = -(\gamma - i\Delta)P + ig\sqrt{n}E + i\Omega S$$

$$\partial_t S = i\Omega^* P$$


sanity check: no atoms

$$g\sqrt{n} = 0$$


$$(\partial_t + c\partial_z)E = 0$$

$$E(z, t) = E(0, t - z/c)$$

undistorted propagation at  $c$



# Two-level medium



$$(\partial_t + c\partial_z)E = ig\sqrt{n}P$$

$$\partial_t P = -(\gamma - i\Delta)P + ig\sqrt{n}E + i\Omega S$$

$$\partial_t S = i\Omega^* P$$

assume: resonant incoming photon, no control

$$\Delta = 0$$


$$\Omega = 0$$

# Two-level medium

$$(\partial_t + c\partial_z)E = ig\sqrt{n}P$$

$$\partial_t P = -(\gamma - i\Delta)P + ig\sqrt{n}E + i\Omega S$$

$$\partial_t S = i\Omega^* P$$



assume: resonant incoming photon, no control

$$\Delta = 0$$

$$\Omega = 0$$

$$(\partial_t + c\partial_z)E = ig\sqrt{n}P$$

$$\partial_t P = -\gamma P + ig\sqrt{n}\hat{E}$$

$$E(z, t) = \int d\omega \tilde{E}(\omega, t) e^{-i\omega t}$$


$$P(z, t) = \int d\omega \tilde{P}(\omega, t) e^{-i\omega t}$$

# Two-level medium

$$(\partial_t + c\partial_z)E = ig\sqrt{n}P$$

$$\partial_t P = -(\gamma - i\Delta)P + ig\sqrt{n}E + i\Omega S$$

$$\partial_t S = i\Omega^* P$$



assume: resonant incoming photon, no control

$$\Delta = 0$$

$$\Omega = 0$$

$$(\partial_t + c\partial_z)E = ig\sqrt{n}P$$

$$\partial_t P = -\gamma P + ig\sqrt{n}\hat{E}$$

$$E(z, t) = \int d\omega \tilde{E}(\omega, t) e^{-i\omega t}$$

$$P(z, t) = \int d\omega \tilde{P}(\omega, t) e^{-i\omega t}$$

$$(-i\omega + c\partial_z)\tilde{E} = ig\sqrt{n}\tilde{P}$$

$$-i\omega\tilde{P} = -\gamma\tilde{P} + ig\sqrt{n}\tilde{E}$$




# Two-level medium

$$(-i\omega + c\partial_z)\tilde{E} = ig\sqrt{n}\tilde{P}$$

$$-i\omega\tilde{P} = -\gamma\tilde{P} + ig\sqrt{n}\tilde{E}$$

$$\tilde{P} = \frac{ig\sqrt{n}}{\gamma - i\omega}\tilde{E}$$



$$c\partial_z\tilde{E} = \left( i\omega - \frac{g^2n}{\gamma - i\omega} \right) \tilde{E}$$

$$\tilde{E}(L, \omega) = \tilde{E}(0, \omega) \exp \left[ i\omega \frac{L}{c} - \frac{g^2nL/c}{\gamma - i\omega} \right]$$


$$\tilde{E}(L, \omega) = \tilde{E}(0, \omega) \exp \left[ i\omega \frac{L}{c} - \frac{d\gamma}{\gamma - i\omega} \right]$$

$$d = \frac{g^2nL}{\gamma c}$$

$$|\tilde{E}(L, \omega)|^2 = |\tilde{E}(0, \omega)|^2 \exp \left[ -\frac{2d}{1 - (\omega/\gamma)^2} \right]$$


# Two-level medium

$$|\tilde{E}(L, \omega)|^2 = |\tilde{E}(0, \omega)|^2 \exp \left[ -\frac{2d}{1 - (\omega/\gamma)^2} \right]$$




- on resonance:  $I_{out} = I_{in} e^{-2d}$
- $2d =$  optical depth
- assume  $d \gg 1$

absorption line



# Electromagnetically induced transparency (EIT)



$$(\partial_t + c\partial_z)E = ig\sqrt{n}P$$

$$\partial_t P = -\gamma P + ig\sqrt{n}E + i\Omega S$$

$$\partial_t S = i\Omega^* P$$

- assume  $\Omega$  real

$$(-i\omega + c\partial_z)\tilde{E} = ig\sqrt{n}\tilde{P}$$

$$-i\omega\tilde{P} = -\gamma\tilde{P} + ig\sqrt{n}\tilde{E} + i\Omega\tilde{S}$$

$$-i\omega\tilde{S} = i\Omega\tilde{P}$$

$$E(z, t) = \int d\omega \tilde{E}(\omega, t) e^{-i\omega t}$$

$$P(z, t) = \int d\omega \tilde{P}(\omega, t) e^{-i\omega t}$$

$$S(z, t) = \int d\omega \tilde{S}(\omega, t) e^{-i\omega t}$$

- at  $\omega = 0$ :  $\tilde{P} = 0$

$$\tilde{S} = -\frac{g\sqrt{n}}{\Omega}\tilde{E}$$


$\partial_z \tilde{E} = 0$  perfect transmission, i.e. no scattering

**Dark-state polariton:** coupled atom-photon excitation

[Fleishhauer & Lukin, 2000, 2002]

- destructive interference

# Electromagnetically induced transparency (EIT)



$$(-i\omega + c\partial_z)\tilde{E} = ig\sqrt{n}\tilde{P}$$


$$-i\omega\tilde{P} = -\gamma\tilde{P} + ig\sqrt{n}\tilde{E} + i\Omega\tilde{S}$$

$$-i\omega\tilde{S} = i\Omega\tilde{P}$$

- near  $\omega = 0$


$$|\tilde{E}(L, \omega)|^2 \approx |\tilde{E}(0, \omega)|^2 \exp\left[-\frac{2d\gamma^2\omega^2}{\Omega^4}\right]$$

homework  
exercise



- EIT transparency window of bandwidth  $\omega_{EIT} \sim \frac{\Omega^2}{\gamma\sqrt{d}}$

# Electromagnetically induced transparency (EIT)



$$(-i\omega + c\partial_z)\tilde{E} = ig\sqrt{n}\tilde{P}$$

$$-i\omega\tilde{P} = -\gamma\tilde{P} + ig\sqrt{n}\tilde{E} + i\Omega\tilde{S}$$

$$-i\omega\tilde{S} = i\Omega\tilde{P}$$

- near  $\omega = 0$

$$\partial_z\tilde{E} \approx i\frac{\omega}{v_g}\tilde{E}$$


$$v_g \approx \frac{\Omega^2}{g^2n}c \ll c$$

homework  
exercise

$$(\partial_t + v_g\partial_z)E = 0$$



reduced group velocity

“slow light”



- pulse compression

# Electromagnetically induced transparency (EIT)




$$\partial_z \tilde{E} \approx i \frac{\omega}{v_g} \tilde{E}$$

$$v_g \approx \frac{\Omega^2}{g^2 n} c \ll c$$

$$(\partial_t + v_g \partial_z) E = 0$$


reduced group velocity

“slow light”



- pulse compression

# Photon storage and retrieval



- dark state polariton


$$|\psi\rangle \sim \int dz f(z - v_g t) \left( \Omega \hat{\mathcal{E}}(z) - g\sqrt{n} \hat{S}^\dagger(z) \right) |0\rangle$$

- while pulse is inside medium, turn  $\Omega$  off

$$|\psi\rangle \sim \int dz f(z) S^\dagger(z) |0\rangle \quad \text{photon stored in "spinwave"}$$

- when turn  $\Omega$  back on, photon is retrieved

$$v_g \approx \frac{\Omega^2}{g^2 n} c \quad \text{reduced group velocity}$$



- pulse compression

# Off-resonant two-level medium


$$(\partial_t + c\partial_z)E = ig\sqrt{n}P$$

$$\partial_t P = -(\gamma - i\Delta)P + ig\sqrt{n}\hat{E} \quad \text{large } \Delta$$

- Fourier transform in time & drop  $\gamma$ :

$$(-i\omega + c\partial_z)\tilde{E} = ig\sqrt{n}\tilde{P}$$

$$-i\omega\tilde{P} \approx i\Delta\tilde{P} + ig\sqrt{n}\tilde{E}$$





# Off-resonant two-level medium

$$(\partial_t + c\partial_z)E = ig\sqrt{n}P$$

$$\partial_t P = -(\gamma - i\Delta)P + ig\sqrt{n}\hat{E} \quad \text{large } \Delta$$

- Fourier transform in time & drop  $\gamma$ :

$$(-i\omega + c\partial_z)\tilde{E} = ig\sqrt{n}\tilde{P}$$

$$-i\omega\tilde{P} \approx i\Delta\tilde{P} + ig\sqrt{n}\tilde{E}$$

- near  $\omega = 0$ :

$$c\partial_z\tilde{E} \approx ig\sqrt{n}\tilde{P}$$


$$0 \approx i\Delta\tilde{P} + ig\sqrt{n}\tilde{E}$$

$$\tilde{P} \approx -\frac{g\sqrt{n}}{\Delta}\tilde{E}$$


$$\partial_z\tilde{E} \approx -i\frac{g^2n}{c\Delta}\tilde{E}$$

$$\tilde{E}(z = L, \omega \approx 0) \approx \tilde{E}(z = 0, \omega \approx 0) \exp\left[-i\frac{d\gamma}{\Delta}\right]$$

- atoms imprint a phase on photon



# EIT with large single-photon detuning



$$(-i\omega + c\partial_z)\tilde{E} = ig\sqrt{n}\tilde{P}$$

$$-i\omega\tilde{P} = -(\gamma - i\Delta)\tilde{P} + ig\sqrt{n}\tilde{E} + i\Omega\tilde{S}$$

large  $\Delta$

$$-i\omega\tilde{S} = i\Omega\tilde{P}$$

• at  $\omega = 0$ :

same as for  $\Delta = 0$

$$\tilde{P} = 0 \quad \tilde{S} = -\frac{g\sqrt{n}}{\Omega}\tilde{E}$$

$$\partial_z\tilde{E} = 0 \quad \text{perfect transmission due to EIT}$$


**Dark-state polariton:** coupled atom-photon excitation  
[Fleishhauer & Lukin, 2000, 2002]

$$(\partial_t + v_g\partial_z)E = 0$$

reduced group velocity

“slow light”

# Medium where photons interact strongly



EIT = electromagnetically induced transparency

Summer school lectures by Browaeys, Hazzard,  
and possibly Kaufman, Bakr, etc...

# Outline

- motivation and basic idea
- E&M field quantization
- propagation of light through atomic ensembles; electromagnetically induced transparency (EIT)
- **Rydberg atoms**
- basic idea revisited
- photon interacting with stationary excitation
  - on resonance: single-photon switch, subtractor
  - off resonance: two-photon quantum gate
- dynamics of multiple photons
  - on resonance: source of single photons
  - off resonance: two-photon gate, bound states, many-body physics
- more applications

electronic levels in atom:



electron wavefunction



nucleus


$n = 1$  —

electronic levels in atom:





$n = 2$  —

$n = 1$  —





electronic levels in atom:






electronic levels in atom:






- large size:  $r \sim n^2$


electronic levels in atom:











Cambridge Hall



Farm Dr

University  
of Maryland  
Parking Lot Q



Computer and Space  
Sciences Building



Astronomy Department



Physical Sciences  
Complex



Joint Quantum Institute



Stadium Dr




Stadium Dr

Stadium Dr

Stadium Dr

Stadium Dr





## Rydberg states


- huge size:  $r \sim n^2$
- huge electric dipole moment
- strong, distant interactions

map on strong, distant photon-photon interactions


# Outline

- motivation and basic idea
- E&M field quantization
- propagation of light through atomic ensembles; electromagnetically induced transparency (EIT)
- Rydberg atoms
- **basic idea revisited**
- photon interacting with stationary excitation
  - on resonance: single-photon switch, subtractor
  - off resonance: two-photon quantum gate
- dynamics of multiple photons
  - on resonance: source of single photons
  - off resonance: two-photon gate, bound states, many-body physics
- more applications

# Medium where photons interact strongly




Map strong **atom-atom interactions** onto strong photon-photon interactions



EIT = electromagnetically induced transparency

# Basic idea


ground-state atoms



- one photon (polariton) drags along a Rydberg excitation

# Basic idea

ground-state atoms



- one photon (polariton) drags along a Rydberg excitation
- another photon drags along a Rydberg excitation
- Rydberg excitations feel strong, distant interactions

⇒ strong, distant photon-photon interactions



# Outline

- motivation and basic idea
- E&M field quantization
- propagation of light through atomic ensembles; electromagnetically induced transparency (EIT)
- Rydberg atoms
- basic idea revisited
- **photon interacting with stationary excitation**
  - on resonance: single-photon switch, subtractor
  - off resonance: two-photon quantum gate
- dynamics of multiple photons
  - on resonance: source of single photons
  - off resonance: two-photon gate, bound states, many-body physics
- more applications